

(iii) *Limitations.* Administer continuously throughout growing period. Do not administer to chickens producing eggs for human consumption. Withdraw 5 days before slaughter. Use as sole source of organic arsenic. Overdosage or the lack of water intake may result in weakness or paralysis of legs.

[46 FR 41040, Aug. 14, 1981, as amended at 46 FR 42448, Aug. 21, 1981; 47 FR 15238, Apr. 9, 1982; 55 FR 8460, Mar. 8, 1990; 57 FR 8577, Mar. 11, 1992; 58 FR 65664, Dec. 16, 1993; 65 FR 10705, Feb. 29, 2000]

§ 520.2089 Roxarsone liquid.

(a) *Specifications.* Each teaspoon (5 milliliters) of solution contains 72 milligrams of roxarsone (3-nitro-4-hydroxyphenylarsonic acid).

(b) *Sponsor.* See No. 046573 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.60 of this chapter.

(d) *Conditions of use in growing chickens and growing turkeys—(1) Amount.* 1 teaspoon (5 milliliters) to each gallon of drinking water (0.002 percent roxarsone).

(2) *Indications for use.* For improved rate of weight gain, improved feed efficiency, and improved pigmentation.

(3) *Limitations.* Administer continuously throughout growing period. Do not administer to chickens producing eggs for human consumption. Withdraw 5 days before slaughter. Use as sole source of organic arsenic. Overdosage or the lack of water intake may result in weakness or paralysis of legs.

[58 FR 65665, Dec. 16, 1993, as amended at 65 FR 10705, Feb. 29, 2000]

§ 520.2095 Sarafloxacin soluble powder.

(a) *Specifications.* Each 145 grams (5.1 ounces) pouch contains sarafloxacin hydrochloride equivalent to 14.5 grams of sarafloxacin base.

(b) *Sponsor.* See No. 000074 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.594 of this chapter.

(d) *Conditions of use.* Used in drinking water as follows:

(1) *Amount.* Chickens—20 to 40 parts per million for 5 consecutive days as the only source of drinking water. Turkeys—30 to 50 parts per million for 5

consecutive days as the only source of drinking water.

(2) *Indications for use.* For control of mortality in growing turkeys and broiler chickens associated with *Escherichia coli* organisms susceptible to sarafloxacin.

(3) *Limitations.* No preslaughter drug withdrawal period is required when the product is used as directed. Use in a manner other than that indicated or with a dose in excess of that recommended may result in drug residues in edible tissues. Do not use in laying hens producing eggs for human consumption. The effects of sarafloxacin on the reproductive function of treated fowl have not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[60 FR 50097, Sept. 28, 1995]

§ 520.2098 Selegiline hydrochloride tablets.

(a) *Specifications.* Each tablet contains either 2, 5, 10, 15, or 30 milligrams of selegiline hydrochloride.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use—Dogs—(1) Dosage.* 1 milligram per kilogram (0.45 milligram per pound) of body weight.

(i) *Indications for use.* For control of clinical signs associated with uncomplicated pituitary-dependent hyperadrenocorticism in dogs.

(ii) *Limitations.* Administer orally once daily. If no improvement in clinical signs or physical examination findings after 2 months of therapy, increase dose to a maximum of 2 milligrams per kilogram once daily. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Dosage.* 0.5 to 1.0 milligram per kilogram of body weight once daily.

(i) *Indications for use.* For the control of clinical signs associated with canine cognitive dysfunction syndrome.

(ii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[62 FR 34632, June 27, 1997; 62 FR 55159, Oct. 23, 1997, as amended at 63 FR 29551, June 1, 1998; 64 FR 2122, Jan. 13, 1999]